

Colour-Octet Contributions to J/ψ Production via Fragmentation at HERA*

BERND A. KNIEHL

Max-Planck-Institut für Physik (Werner-Heisenberg-Institut),
Föhringer Ring 6, 80805 Munich, Germany

Abstract

We study J/ψ photoproduction via fragmentation at next-to-leading order in the QCD-improved parton model, using the nonrelativistic factorization formalism proposed by Bodwin, Braaten, and Lepage. We predict that measurements of J/ψ photoproduction at DESY HERA should show a distinctive excess over the expectation based on the colour-singlet model at small values of the inelasticity variable z .

Since its discovery 25 years ago, the J/ψ meson has provided a useful laboratory for quantitatively testing quantum chromodynamics (QCD) and, in particular, the interplay of perturbative and nonperturbative phenomena. Recently, the cross section of J/ψ inclusive production measured at the Fermilab Tevatron turned out to be more than one order of magnitude in excess of what used to be the best theoretical prediction, based on the colour-singlet model (CSM). As a solution to this puzzle, Bodwin, Braaten, and Lepage [1] proposed the existence of so-called colour-octet processes to fill the gap. The idea is that $c\bar{c}$ pairs are produced at short distances in colour-octet states and subsequently evolve into physical (colour-singlet) charmonia by the nonperturbative emission of soft gluons. The underlying theoretical framework is provided by nonrelativistic QCD (NRQCD) endowed with a particular factorization theorem, which implies a separation of short-distance coefficients, which are amenable to perturbative QCD, from long-distance matrix elements, which must be extracted from experiment. This formalism involves a double expansion in the strong coupling constant α_s and the relative velocity v of the bound c quarks, and takes the complete structure of the charmonium Fock space into account.

In order to convincingly establish the phenomenological significance of the colour-octet mechanism, it is indispensable to identify it in other kinds of high-energy experiments as well. In this presentation, we briefly report on a next-to-leading-order (NLO) analysis

*To appear in *Proceedings of the International Europhysics Conference on High Energy Physics*, Jerusalem, Israel, 19-26 August 1997.

[2] of inelastic J/ψ photoproduction via fragmentation at HERA. We study direct and resolved photoproduction of prompt J/ψ mesons and χ_{cJ} mesons ($J = 0, 1, 2$) that radiatively decay to $J/\psi + \gamma$, taking into account the formation of both colour-singlet and colour-octet $c\bar{c}$ states. The dominant J/ψ (χ_{cJ}) Fock states are $[\underline{1}, ^3S_1]$ and $[\underline{8}, ^3S_1]$ ($[\underline{1}, ^3P_J]$ and $[\underline{8}, ^3S_1]$), respectively. For comparison, we also consider γg fusion in the CSM.

From Fig. 1a, we observe that the fragmentation mechanism vastly dominates inelastic J/ψ photoproduction in the lower range of the inelasticity variable $z = p_p \cdot p_{J/\psi} / p_p \cdot p_\gamma$. For a minimum- p_T cut of 4 GeV (8 GeV), its contribution exceeds the one due to γg fusion for $z < 0.4$ (0.75), by factors of about 4 and 200 (20 and 700) at $z = 0.25$ and 0.05, respectively. The bulk of the fragmentation contribution is induced by resolved photons. In Fig. 1b, the ZEUS $J/\psi \rightarrow \mu^+\mu^-$ data [3] collected in the interval $0.3 < z < 0.8$ are compared with the respective fragmentation and γg -fusion predictions. We have divided the experimental data points by the estimated extrapolation factor 1.07 which was included in Ref. [3] to account for the unmeasured contribution from $0 < z < 0.3$. In fact, our combined analysis of fragmentation and γg fusion suggests that, at $p_T = 5$ GeV, this factor should be as large as 3.4. The corresponding value for $0.05 < z < 0.3$ is 2.6.




Figure 1: (a) $d\sigma/dz$ and (b) $d\sigma/dp_T^2$ for inelastic J/ψ photoproduction at HERA.

In conclusion, the cross section of inelastic J/ψ photoproduction in ep collisions at low to intermediate z and large p_T , where fragmentation production is dominant, is very sensitive to the colour-octet matrix element $\langle 0 | \mathcal{O}^{J/\psi}[\underline{8}, ^3S_1] | 0 \rangle$. By contrast, γg fusion, which is only relevant in the upper z range and for $p_T \lesssim M_{J/\psi}$, probes $\langle 0 | \mathcal{O}^{J/\psi}[\underline{8}, ^1S_0] | 0 \rangle$ and $\langle 0 | \mathcal{O}^{J/\psi}[\underline{8}, ^3P_J] | 0 \rangle$. We propose to accordingly extend previous measurements of this cross section at HERA, in order to obtain an independent, nontrivial check of the Tevatron colour-octet charmonium puzzle.

Acknowledgements. I would like to thank Gustav Kramer for his collaboration on the work presented here.

References

- [1] G.T. Bodwin, E. Braaten, and G.P. Lepage, Phys. Rev. D 51 (1995) 1125; 55 (1997) 5855 (E).
- [2] B.A. Kniehl and G. Kramer, Phys. Rev. D 56 (1997) 5820; Phys. Lett. B 413 (1997) 416.
- [3] ZEUS Collaboration, J. Breitweg et al., Contributed Paper No. pa02–047 to the 28th International Conference on High Energy Physics, Warsaw, Poland, 25–31 July 1996.